

260 mm

10 mm



**DISPERSIBLE GRANULES**

<b>ACTIVE INGREDIENT</b>	<b>BY WEIGHT</b>
Mancozeb: A coordination product of zinc ion and manganese ethylenebisdithiocarbamate	75.0%
in which the ingredients are:	
Manganese++	15.0%
Zinc++	1.9%
Ethylenebisdithiocarbamate ion (C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> S <sub>4</sub> ) --	58.1%
<b>OTHER INGREDIENTS</b>	<b>25.0%</b>
<b>TOTAL</b>	<b>100.0%</b>

Contains 0.75 Pound of Mancozeb Per Pound of Product

EPA Reg. No. 70506-234

EPA Est. No. 70506-COL-001

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**



See attached booklet on back for additional Precautionary Statements and complete Directions For Use.



**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300.**

**Net Weight: 6 Pounds**

2180181-A (0223)  
ESL010722-10041-022422



Made in Colombia  
**UPL NA Inc.** • 630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406 U.S.A. • 1-800-438-6071



10 mm

50 mm

100 mm

50 mm

10 mm

880 mm

385 mm

ESL010722-10041-022422  
2180181-A (0223)

UPL, the UPL logo, MANZATE and PRO-STICK are trademarks of a UPL Corporation Limited Group Company.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage or disposal.  
**PESTICIDE STORAGE:** Important-Never allow MANZATE PRO-STICK to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of MANZATE PRO-STICK as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed in a secure dry area.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**DIRECTIONS FOR USE**  
This product is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water by disposing of equipment washwater or rinsate. Cover or incorporate spilled treated seed.

**ENVIRONMENTAL HAZARDS**  
Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, clothing or eyes. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION  
PRECAUTIONARY STATEMENTS**



<b>FIRST AID</b>	
<b>IF ON SKIN OR CLOTHING</b>	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
<b>IF IN EYES</b>	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<b>IF INHALED</b>	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably by mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<b>IF SWALLOWED</b>	• Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical assistance, call Rocky Mountain Poison and Drug Safety at 1-866-673-6671.	

10 mm

240 mm



GROUP M3 FUNGICIDE

# MANZATE<sup>®</sup> PRO-STICK<sup>™</sup>

FUNGICIDE

See inside for complete Precautionary Statements, First Aid and Directions For Use.

FUNGICIDE

Made in Colombia

UPL NA Inc. • 630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406 U.S.A. • 1-800-438-6071



# MANZATE<sup>®</sup> PRO-STICK<sup>™</sup>

## FUNGICIDE

### DISPERSIBLE GRANULES

#### ACTIVE INGREDIENT

BY WEIGHT

Mancozeb: A coordination product of zinc ion and manganese ethylenebisdithiocarbamate..... 75.0%

in which the ingredients are:

Manganese++..... 15.0%

Zinc++..... 1.9%

Ethylenebisdithiocarbamate ion (C<sub>4</sub>H<sub>6</sub>N<sub>2</sub>S<sub>4</sub>) --..... 58.1%

OTHER INGREDIENTS..... 25.0%

TOTAL..... 100.0%

Contains 0.75 Pound of Mancozeb Per Pound of Product

EPA Reg. No. 70506-234

EPA Est. No. 70506-COL-001

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300.

## FIRST AID

<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably by mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical assistance, call Rocky Mountain Poison and Drug Safety at 1-866-673-6671.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300.**

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **CAUTION**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, clothing or eyes. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are nitrile rubber, natural rubber, or butyl rubber.

**Mixers, loaders, applicators and other handlers must wear:**

- Long-sleeved shirt
- Long pants
- Shoes and socks
- Chemical-resistant gloves, made of any waterproof material (except pilots, groundboom applicators, and airblast applicators)

Aerial applications of MANZATE® PRO-STICK™ Fungicide on broccoli, cabbage, lettuce (leaf and head), and peppers requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a minimum of a NIOSH-approved particulate filtering facepiece respirator with R or P filter, OR a NIOSH-approved elastomeric particulate respirator with any R or P filter.

**In addition mixers/loaders supporting aerial applications to walnuts must wear:** a minimum of a NIOSH-approved particulate filtering facepiece respirator with R or P filter, NIOSH-approved elastomeric particulate respirator with any R or P filter.

See engineering controls for additional requirements.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

## ENGINEERING CONTROL STATEMENTS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for [40 CFR Part 170.240 (d)(4-6)].

Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR Part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## **USER SAFETY RECOMMENDATIONS**

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water by disposing of equipment washwater or rinsate. Cover or incorporate spilled treated seed.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material

## **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Commercial seed treatments and applications to lawn grasses, golf courses, industrial (office park), and municipal lawns are not within the scope of the Worker Protection Standard.

Do not enter or allow others to enter treated areas until sprays have dried.

MANZATE PRO-STICK, a dispersible granule containing mancozeb, is labeled for use as a spray for the control of many important plant diseases.

## **APPLICATION INSTRUCTIONS**

AS A SPRAY (Ground or Aerial Equipment) - Apply MANZATE PRO-STICK at the rate shown; use sufficient water to provide thorough coverage: use 20 to 100 gallons per acre for ground equipment and no less than 2 gallons per acre for aircraft. Add MANZATE PRO-STICK slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension. A spreader-sticker spray adjuvant may be used with this product if needed; contact your local product distributor or UPL NA Inc. representative for specific recommendations. If tank mixed, follow more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on tank mixing.

## **RESISTANCE MANAGEMENT**

MANZATE PRO-STICK Fungicide is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. MANZATE PRO-STICK is recommended for use in program which are compatible with the principles of Integrated Pest Management (IPM). MANZATE PRO-STICK is effective for strategic use in programs that attempt to minimize disease resistance to fungicides.

Some other fungicides which are at risk from disease resistance exhibit a single-site mode of fungicidal action. MANZATE PRO-STICK, with a multi-site mode of action, may be used in an Integrated Pest Management (IPM) program to delay or prevent the development of resistance to single-site fungicides.

Consult with your private crop consultant or Federal or State Cooperative Extension Service representatives for guidance on the proper use of MANZATE PRO-STICK in programs which seek to minimize the occurrence of disease resistance to other fungicides.

## **MAXIMUM SEASONAL POUNDAGE WHEN USED IN CONJUNCTION WITH ONE OR MORE OTHER SPECIFIED ETHYLENE BISDITHIOCARBAMATE (EBDC) PRODUCTS**

### **Foliar Applications**

#### **Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season**

If more than one product containing an EBDC active ingredient (mancozeb or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

#### **Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season**

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

### **Seed Treatment**

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application for seed treatment may be made on crops which have registered seed treatment uses.

### **CHEMIGATION**

Apply MANZATE PRO-STICK Fungicide only through sprinkler systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems. Do not apply MANZATE PRO-STICK through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.



### **Specific Instructions for Public Water Systems:**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

### **Specific Instructions for Sprinkler Irrigation Systems:**

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of MANZATE PRO-STICK as a continuous injection. In non-moving systems inject MANZATE PRO-STICK for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of MANZATE PRO-STICK needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all MANZATE PRO-STICK is flushed from system.

## **SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

### Wind Speed

Do not apply at wind speeds greater than 15 mph.

### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of mancozeb. Where states have more stringent regulations, they must be observed.

### Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

*Additional requirements for aerial applications:*

1. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
2. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
3. When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

*Additional requirements for ground boom application:*

1. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

## **Tank Mixing Procedures**

- When two pesticides are tank mixed, the more restrictive label conditions apply.
- Do not tank mix with any product which contains a restriction on tank mixing.

Slowly pour into spray tank as long as it is being filled while agitation is maintained or thoroughly premix in a nurse tank for concentrate or aircraft sprayers. Add other co-applied fungicides, insecticides, growth regulators, micronutrients after MANZATE PRO-STICK has been placed into suspension. When tank mixing with other pesticides, observe the more restrictive label limitations, restrictions, and precautions.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When preparing spray solutions for use in a hand sprayer, premix as a slurry in a small container, and then add to sprayer containing 1/3 to 1/2 the desired final water volume.

## **Compatibility**

MANZATE PRO-STICK is compatible with most commonly used agricultural fungicides, insecticides and growth regulators. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

## CROP USE TABLE AND APPLICATION DIRECTIONS

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Almond	Anthracnose Blossom Blight <i>(Monilinia spp.)</i> Rust Scab Shothole <i>(Stigmina spp.)</i>	6.4 lbs/A (4.8 lbs ai/A)	Begin application at dormant to popcorn stage, full bloom or petal fall. Reapply every 7 to 10 days if bloom is staggered and weather is rainy.  For increased control of Bacterial Spot, tank mix MANZATE PRO-STICK with copper or an EPA approved bactericide registered for use on almonds.	Do not apply more than 19.2 lbs of product (14.4 lbs of active ingredient) per acre per use season. Maximum number of applications per season is 3.  Do not make last application later than 5 weeks after petal fall.  Do not graze livestock in treated area. Do not use less than 10 gallons of spray volume per acre if aerially applied.  Minimum retreatment interval is 7 days.  Do not apply this product with a U-boom device.

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Asparagus	Cercospora Leaf Spot Rust	2 lbs/A (1.50 lbs ai/A)	Start applications when disease first appears and repeat at 10 day intervals.	Apply only on asparagus ferns after spears have been harvested. Do not apply more than 8 lbs of product (6 lbs of active ingredient) per acre per season. Maximum number of applications per season is 4.  Preharvest Interval (PHI) for California and Arizona: 120 days.  Preharvest Interval (PHI) for all other states: 180 days
Asparagus Crown (Planting Stock)	Crown Rot	1.0 lb/100 gals (0.75 lbs ai/100 gals)	Dip clean, loosely packed crowns into continuously agitated fungicide suspension for 5 minutes. Drain and plant as soon as possible. Wash dirty crowns before dip treatment. Replace suspension in clean tank when discolored by soil.	

*(continued)*

<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)</b>	<b>APPLICATION DIRECTIONS</b>	<b>RESTRICTIONS</b>
Banana (Including Plantain)	Sigatoka	2 to 3 lbs/A (1.50 to 2.25 lbs ai/A)	Apply when leaves first appear and repeat every 14 to 21 days or as required. Use sufficient water to provide adequate coverage.	Do not apply more than 30 lbs of product (22.5 lbs of active ingredient) per acre per growing cycle. Maximum number of applications per season is 10 at the highest use rate; 15 at the lowest use rate. Preharvest Interval (PHI): 0 days
Cereals Barley Oat Rye Wheat (Including Triticale)	Helminthosporium Leaf Spot Leaf Rust Septoria Glume Blotch Septoria Leaf Spot Tan Spot	2 lbs/A (1.50 lbs ai/A)	For leaf and stem diseases, start application at onset of disease or when plants are in the tillering to jointing stage and repeat at 7 to 10 day intervals. For Fusarium head blight in barley the optimum timing for suppression is when barley heads on the main stem are fully emerged which is Feekes growth stage 10.51. For Fusarium head blight in wheat the optimum timing for suppression is at early flower which is Feekes growth stage 10.51. Good coverage of crop heads is required.	Do not apply more than 6 lbs of product (4.5 lbs of active ingredient) per acre per crop. Maximum number of applications per season is 3. Do not graze livestock in treated areas prior to harvest. Preharvest Interval (PHI): 26 days

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Broccoli	Alternaria Leaf Spot Downy Mildew	1.6 to 2.1 lbs/A (1.2 to 1.575 lbs ai/A)	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 12.8 lbs of product (9.6 lbs of active ingredient) per acre per season. Maximum number of applications per season is 6 at the highest use rate; 8 at the lowest use rate. Minimum retreatment interval is 7 days. Do not apply this product with a U-boom device. Aerial application of MANZATE PRO-STICK on broccoli, requires that occupational handlers performing mixing/loading operational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a minimum of a NIOSH-approved particulate filtering facepiece respirator with an N, R, or P filter, NIOSH approval prefix TC-84A. Preharvest Interval (PHI): 7 days

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Cabbage	Alternaria Leaf Spot Downy Mildew	1.6 to 2.1 lbs/A (1.2 to 1.575 lbs ai/A)	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 12.8 lbs of product (9.6 lbs of active ingredient) per acre per season. Maximum number of applications per season is 6 at the highest use rate; 8 at the lowest use rate. Minimum retreatment interval is 7 days. Do not apply this product with a U-boom device. Aerial application of MANZATE PRO-STICK on cabbage, requires that occupational handlers performing mixing/loading operational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a minimum of a NIOSH-approved particulate filtering facepiece respirator with an N, R, or P filter, NIOSH approval prefix TC-84A. Preharvest Interval (PHI): 7 days

*(continued)*



CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Caprifig (Non-Food Use)	Endosepsis ( <i>Fusarium</i> ) Mold	4 lbs/100 gals (3.0 lbs ai/100 gals)	Prepare mamme figs by making a shallow cut through the eye and then hand dividing to avoid wasp injury. Submerge mamme figs in the continuously agitated suspension for at least 15 minutes. Drain before placement in trees. Use fresh dipping suspension after treating 4 to 5 batches of figs.	

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Corn (Sweet Corn for Fresh Use or Processing; Popcorn; and Sweet Corn for Seed Production, including Hybrid Seed)	Common Rust Helminthosporium Leaf Blight Gray Leaf Spot	1.5 lbs/A (1.125 lbs ai/A)	Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 4 to 7 day intervals.	Sweet Corn: Do not apply more than 22.5 lbs of product (17 lbs of active ingredient) per acre per crop east of the Mississippi River (including Arkansas and Louisiana). Maximum number of applications per season is 15. Do not apply more than 7.5 lbs of product (5.6 lbs of active ingredient) per acre per crop west of the Mississippi River (excluding Arkansas and Louisiana). Maximum number of applications per season is 5. Do not apply to sweet corn in home gardens. Preharvest interval (PHI): 7 days
(Field and Field Corn for Hybrid Seed Production)				Field Corn: Do not apply more than 15 lbs of product (11.25 lbs of active ingredient) per acre per crop. Maximum number of applications per season is 10. Preharvest interval (PHI): 40 days

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Cranberry	Fruit Rot	3 to 6 lbs/A (2.25 to 4.5 lbs ai/A)	Start applications at early-bloom and repeat at 7 to 10 day intervals.	Do not apply more than 18 lbs of product (13.5 lbs of active ingredient) per acre per season.  Maximum number of applications per season is 3 at the highest use rate; 6 at the lowest use rate.  Preharvest interval (PHI): 30 days
Cucurbit Crop Group Chayote Chinese wax gourd Citron melon Cucumber Gherkin Gourd, edible <i>Momordica</i> spp. Muskmelon Pumpkin Squash, summer Squash, winter Watermelon	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight Scab	2 to 3 lbs/A (1.50 to 2.25 lbs ai/A)	Start applications when the plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.  For aerial applications, the minimum spray volume is 2 gallons per acre.  Some cantaloupe varieties (i.e. Harvest Queen, Gold Star, Super Star, Sweet and Early, and Saticoy) are sensitive to MANZATE PRO-STICK Fungicide.  Consult State Cooperative Extension Service Specialist prior to use.	Do not apply more than 25.6 lbs of product (19.2 lbs of active ingredient) per acre per year.  Maximum number of applications per year is 8 at the highest use rate; 12 at the lowest use rate.  Preharvest Interval (PHI): 5 days

<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)</b>	<b>APPLICATION DIRECTIONS</b>	<b>RESTRICTIONS</b>
Fennel	Early Blight Late Blight	2 lbs/A (1.5 lbs ai/A)	Begin in plant beds at emergence. Repeat at 7 to 10 day intervals.	Do not apply more than 16 lbs of product (12 lbs of active ingredient) per acre per crop. Maximum number of applications per year is 8. Do not graze livestock in treated areas. Preharvest Interval (PHI): 14 days
Ginseng	Alternaria Blight	2 lbs/A (1.5 lbs ai/A)	Start applications when disease first threatens and repeat every 7 to 10 days as needed. In Wisconsin, apply with ground equipment and a minimum of 80 gallons of water per acre.	Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per year. Maximum number of applications per year is 12. Preharvest Interval (PHI): 30 days

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Grape (East of the Rocky Mountains)	Black Rot Bunch Rot Deadarm Downy Mildew	1.5 to 4 lbs/A (1.125 to 3.0 lbs ai/A)	Apply in sufficient water to provide thorough coverage starting when new shoots are 1/2 to 1 1/2 inches long. Repeat when shoots are 3 to 5 inches long, when shoots are 8 to 10 inches long, and then at 7 to 10 day intervals until fruit is set. For late season control of black rot, deadarm and downy mildew, the use of other approved fungicides is suggested.	Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per season. Maximum number of applications per season is 6 at the highest use rate; 16 at the lowest use rate. Preharvest Interval (PHI): 66 days
Grape (West of the Rocky Mountains)	Black Rot Bunch Rot Deadarm Downy Mildew	1.5 to 2.5 lbs/A (1.125 to 1.875 lbs/A)	Apply in sufficient water to provide thorough coverage starting when new shoots are 1/2 to 1 1/2 inches long. Repeat when shoots are 3 to 5 inches long, when shoots are 8 to 10 inches long, and then at 7 to 10 day intervals until fruit is set. For late season control of black rot, deadarm and downy mildew, the use of other approved fungicides is suggested.	Do not apply more than 7.5 lbs of product (5.6 lbs of active ingredient) per acre per season. Maximum number of applications per season is 3 at the highest use rate; 5 at the lowest use rate. Preharvest Interval (PHI) for California: No application after bloom. Preharvest Interval (PHI) all other states: 66 days

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Lettuce (Head, Leaf)	Anthracnose Downy Mildew	1.6 to 2.1 lbs/A (1.2 to 1.575 lbs ai/A)	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Remove residues from head lettuce by stripping and trimming. In California, do not apply more than 8.5 lbs of product (6.4 lbs of active ingredient) per acre per crop. Maximum number of applications in California per season is 4 at the highest use rate; 5 at the lowest use rate. In states other than California, do not apply more than 12.8 lbs of product (9.6 lbs of active ingredient) per acre per crop. For all other states, the maximum number of applications per season is 6 at the highest use rate; 8 at the lowest use rate. Minimum retreatment interval is 7 days. Do not apply this product with a U-boom device.

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Lettuce (Head, Leaf)	Anthracnose Downy Mildew	1.6 to 2.1 lbs/A (1.2 to 1.575 lbs ai/A)	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Aerial application of MANZATE PRO-STICK on lettuce (leaf and head), requires that occupational handlers performing mixing/loading operational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a minimum of a NIOSH-approved particulate filtering facepiece respirator with an N, R, or P filter, NIOSH approval prefix TC-84A.  Preharvest Interval (PHI) in California: 14 days Preharvest Interval (PHI) in all other states: 10 days

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Melon Cantaloupe Casaba Crenshaw Honeydew Watermelon	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight	2 to 3 lbs/A (1.50 to 2.25 lbs ai/A)	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals.  Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces. Some varieties are sensitive to MANZATE PRO-STICK Fungicide.  Consult State Cooperative Extension Service Specialist prior to use.	Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per season.  Maximum number of applications per season is 8 at the highest use rate; 12 at the lowest use rate.  Preharvest Interval (PHI): 5 days

*(continued)*



CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS ai/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Onion (Dry Bulb) Garlic Shallot	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	3 lbs/A (2.25 lbs ai/A)	Follow a protective spray schedule starting when diseases are first reported in the area and repeat at 7 day intervals throughout the season.	Do not apply to exposed bulbs. Do not apply more than 30 lbs of product (22.5 lbs of active ingredient) per acre per crop. Maximum number of applications per season is 10. Preharvest Interval (PHI): 7 days
	Smut (furrow drench)	3 lbs/A (2.25 lbs ai/A)	Apply 3 lbs per acre as a furrow drench at time of planting onion seeds. Use 75 to 125 gallons of water per acre.	Do not use more than 3 lbs of product (2.25 lbs of active ingredient) per acre (29,000 linear feet of furrow) with an 18 inch row spacing.

*(continued)*

<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS ai/A)</b>	<b>APPLICATION DIRECTIONS</b>	<b>RESTRICTIONS</b>
Papaya	Anthracnose (Colletotricum) Black Spot (Cercospora) Fruit Rot Phytophthora	2.0 to 2.5 lbs/A (minimum 50 gals of water per acre) (1.50 to 1.875 lbs ai/A)	Begin at flowering; treat central column crown, blossom area and developing fruit. Repeat at 14 to 21 day intervals.	Do not use more than 37 lbs of product (28 lbs of active ingredient) per acre per year. Maximum number of applications per year is 14 at the highest use rate; 18 at the lowest use rate. Preharvest Interval (PHI): 0 days
Peanut	Ascochyta Web Blotch Cercospora Leaf Spot Rust	1 to 2 lbs/A (0.75 to 1.50 lbs ai/A)	Start application when disease first appears or is reported in the area. Repeat sprays at 7 to 14 day intervals. Reduce sprays to a 7 day interval during humid weather or under severe disease pressure.	Do not use more than 16 lbs of product (12 lbs of active ingredient) per acre per crop. Maximum number of applications per season is 8 at the highest use rate; 16 at the lowest use rate. Do not feed treated vines to livestock. Preharvest Interval (PHI): 14 days

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Pepper	Anthracnose Bacterial Spot ( <i>Xanthomonas axonopodis</i> pv. <i>vesicatoria</i> ) Cercospora Leaf Spot (Frogeye Spot) Phytophthora Blight Ripe Rot	West of the Mississippi: 1.6 to 2.1 lbs/A (1.2 to 1.575 lbs ai/A)	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 12.8 lbs of product (9.6 lbs of active ingredient) per acre per season. Maximum number of applications per season is 6 at the highest use rate; 8 at the lowest use rate. Minimum retreatment interval is 7 days. Do not apply this product with a U-boom device. Aerial application of MANZATE PRO-STICK on peppers, requires that occupational handlers performing mixing/loading operational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a minimum of a NIOSH-approved particulate filtering facepiece respirator with an N, R, or P filter, NIOSH approval prefix TC-84A. Preharvest Interval (PHI): 7 days

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Pepper	Anthracnose Bacterial Spot ( <i>Xanthomonas axonopodis</i> pv. <i>vesicatoria</i> ) Cercospora Leaf Spot (Frogeye Spot) Phytophthora Blight Ripe Rot	East of the Mississippi: 1.6 to 3.2 lbs/A (1.2 to 2.4 lbs ai/A)	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 19.2 lbs of product (14.4 lbs of active ingredient) per acre per season). Maximum number of applications per season is 6 at the highest use rate; 12 at the lowest use rate.  Minimum retreatment interval is 7 days. Do not apply this product with a U-boom device.  Aerial application of MANZATE PRO-STICK on peppers, requires that occupational handlers performing mixing/loading operational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a minimum of a NIOSH-approved particulate filtering facepiece respirator with an N, R, or P filter, NIOSH approval prefix TC-84A. Preharvest Interval (PHI): 7 days

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Pome Fruit Apple Pear Crabapple Quince	Rusts Scab Fabrea Leaf Spot	6.0 lbs/A (4.5 lbs ai/A) Maximum per acre use rate based on thorough coverage dilute sprays. Use 50 gal minimum per acre. Consult State Extension Service if necessary to adjust for variable tree size.	<b>Pre-Bloom/Bloom Use:</b> Begin application at 1/4 to 1/2 inch green tip and continue on a 7 to 10 day schedule through bloom. Use either the "Pre-Bloom/Bloom" or "Extended Application" schedule.	Do not apply after petal fall for these diseases. Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per year. Maximum number of applications per season is 3 when using to control Pear Psylla Nymphs or 4 applications when using the higher rates as a pre-bloom/bloom use. If Extended Application Schedule is used, maximum number of applications per season is 8. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program. Preharvest Interval (PHI): 77 days <b>DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.</b>

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Pome Fruit Apple Pear Crabapple Quince	Rusts Scab Fabrea Leaf Spot	3.0 lbs/A (2.25 lbs ai/A) Maximum per acre use rate based on thorough coverage dilute sprays. Use 50 gal minimum per acre. Consult State Extension Service if necessary to adjust for variable tree size.	<u><b>Extended Application Schedule or for Use in Tank Mixtures:</b></u> For implementation of IPM programs, applications based on tree-row volume, or for use as a resistance management tool: begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7 to 10 day schedule through the second cover spray. Use either the "Pre-Bloom/Bloom" or "Extended Application" schedule.	Do not apply after petal fall for these diseases. Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per year. Maximum number of applications per season is 3 when using to control Pear Psylla Nymphs or 4 applications when using the higher rates as a pre-bloom/bloom use. If Extended Application Schedule is used, maximum number of applications per season is 8. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program. Preharvest Interval (PHI): 77 days <b>DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.</b>

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Potato	Black Dot Early Blight Late Blight <b>Disease Suppression:</b> Botrytis	1 to 2 lbs/A (0.75 to 1.50 lbs ai/A)	Begin applications when plants are 4 to 6 inches high by applying 1 lb per acre. As the vines increase in size, apply 1.5 to 2 lbs per acre at intervals of 5 to 10 days or 1 lb per acre at 3 to 5 day intervals.  It is recommended that this product be used in an Integrated Pest Management Program.	Do not apply more than 15 lbs of product (11.2 lbs of active ingredient) per acre per crop.  Maximum number of applications per season is 7 at the highest use rate; 15 at the lowest use rate.  Vine-kill should occur 14 days before harvest.  Preharvest Interval (PHI) in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT, WI: 3 days Preharvest Interval (PHI) in all other states: 14 days

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Potato (Seedpiece Treatment)	Fusarium Decay Seedborne Common Scab	1.25 lbs per 50 gal (0.94 lb ai/50 gal)	Dip whole or cut potato tubers in 1.25 lbs MANZATE PRO-STICK Fungicide per 50 gallons of water. Place treated tubers in a clean container following treatment and plant as soon as possible. Spread treated seedpieces in a cool place if held before planting.	Do not use treated seed potatoes for food or feed purposes. When opening this bag or loading/pouring the treated seed-pieces, users must wear long-sleeved shirt, long pants, shoes, socks, chemical-resistant gloves, and a minimum of a NIOSH-approved particulate filtering facepiece respirator with a an N, R, or P filter, NIOSH-approved prefix TC-84A. After the seed pieces have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: Once the seed pieces are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the soil/media subsurface. <b>SEE ADDITIONAL SEED TREATMENT LANGUAGE FOR SEED BAG LABEL REQUIREMENTS.</b>



<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)</b>	<b>APPLICATION DIRECTIONS</b>	<b>RESTRICTIONS</b>
Squash (Summer Squash, Including Edible Gourd)	Downy Mildew	2 to 3 lbs/A (1.50 to 2.25 lbs ai/A)	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals.  Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.	Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per crop.  Maximum number of applications per season is 8 at the highest use rate; 12 at the lowest use rate.  Preharvest Interval (PHI): 5 days
Sugar Beet	Cercospora Leaf Spot	1.5 to 2 lbs/A (1.125 to 1.50 lbs ai/A)	Begin when disease first threatens. Repeat at 7 to 10 day intervals.	Do not apply more than 14 lbs of product (10.5 lbs of active ingredient) per acre per crop.  Maximum number of applications per season is 7 at the highest use rate; 9 at the lowest use rate.  Do not feed treated sugar beet tops to livestock.  Preharvest Interval (PHI): 14 days

*(continued)*

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS ai/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Tomato (East of the Mississippi River)	Anthracnose Early Blight Gray Leaf Mold Gray Leaf Spot Late Blight Septoria Leaf Spot	0.75 to 3.0 lbs/A (0.56 to 2.25 lbs ai/A)	Start application when seedlings emerge or transplants are set. When using use rate of 0.75 - 1.5 lbs/A, repeat at 3 to 7 day intervals throughout the season. When using use rate of 1.6 - 3.0 lbs/A, repeat at 7 to 10 day intervals throughout the season.	Do not apply more than 22.4 lbs of product (16.8 lbs of active ingredient) per acre per crop. Maximum number of applications is 7 at the highest use rate; 29 at the lowest use rate. Preharvest Interval (PHI): 5 days
	Bacterial Speck and Spot	1.5 to 3 lbs/A (1.125 to 2.25 lbs ai/A)	Start application when seedlings emerge or transplants are set. Repeat at 7 to 10 day intervals throughout the season. For Bacterial Speck and Spot use a full rate of a fixed copper fungicide in a tank mix combination with a half to full rate of MANZATE PRO-STICK. Follow the application interval on the copper fungicide label.	

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS ai/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Tomato (West of the Mississippi River)	Anthracnose Early Blight Gray Leaf Mold Gray Leaf Spot Late Blight Septoria Leaf Spot	0.75 to 2.0 lbs/A (0.56 to 1.50 lbs ai/A)	Start application when seedlings emerge or transplants are set. When using use rate of 0.75 - 1.0 lbs/A, repeat at 3 to 7 day intervals throughout the season. When using use rate of 1.5 - 2.0 lbs/A, repeat at 7 to 10 day intervals throughout the season.	Do not apply more than 8.5 lbs of product (6.4 lbs of active ingredient) per acre per crop. Maximum number of application is 4 at the highest use rate; 11 at the lowest use rate. Preharvest Interval (PHI): 5 days
	Bacterial Speck and Spot	1.5 to 2 lbs/A (1.125 to 1.50 lbs ai/A)	Start application when seedlings emerge or transplants are set. For Bacterial Speck and Spot use a full rate of a fixed copper fungicide in a tank mix combination with a half to full rate of MANZATE PRO-STICK. Follow the application interval on the copper fungicide label. Repeat at 7 to 10 day intervals throughout the season.	

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS ai/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Small Tropical Fruits Limited to: Sugar apple Cherimoya Atemoya Custard apple Sweetsop	Anthracnose	2 to 2.5 lbs/A (1.50 to 1.875 lbs ai/A)	Begin applications at flowering and continue at a 7 day retreatment interval.  Applications made with aerial equipment must be made in a minimum spray volume of 10 gal/acre.	Do not apply more than 35 lbs of product (26.25 lbs of active ingredient) per acre per year.  Maximum number of applications per year is 14 at the highest use rate; 17 at the lowest use rate.  Preharvest Interval (PHI): 0 days
Large Tropical Fruits Limited to: Mango Star apple (caimito) Canistel Mamey sapote Sapodilla White sapote	Anthracnose Phytophthora Fruit Rot Black Spot ( <i>Cercospora</i> )	2 to 2.5 lbs/A (1.50 to 1.875 lbs ai/A)	Start applications at flowering and continue at 14 to 21 day intervals. Direct spray to crown and blossom area. Use 20 to 100 gallons water per acre.	Do not apply more than 37.3 lbs of product (28 lbs of active ingredient) per acre per year.  Maximum number of applications per year is 15 at the highest use rate; 18 at the lowest use rate.  Preharvest Interval (PHI): 0 days

(continued)

CROP	DISEASES CONTROLLED	RATE OF MANZATE PRO-STICK PER APPLICATION LBS PRODUCT/ACRE (LBS AI/A)	APPLICATION DIRECTIONS	RESTRICTIONS
Walnut	Walnut Blight <i>(Xanthomonas xampestris</i> pv. <i>Juglandis)</i>	2.4 lbs/A (1.80 lbs ai/A)	Apply by ground in a minimum of 100 gallons water per acre, or by air in a minimum of 10 gallons water per acre. Begin application at early pre-bloom prior to or when catkins are partially expanded.  Make additional application during bloom and early nutlet stage at 7 to 10 day intervals, or as needed if frequent rainfall occurs.	Do not apply through any irrigation system.  Do not apply more than 24 lbs of product (18 lbs of active ingredient) per acre per use season. Maximum number of applications per season is 10.  For aerial applications only: Mixer/loaders must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with an N, R, or P filter, NIOSH approval prefix TC-84A.  This product must be tank mixed with a fixed copper product (such as Cuprofix Ultra 40 Disperss, EPA Reg. No. 70506-201) or another bactericide which is registered for use on walnuts.  Preharvest Interval (PHI): 75 days

## GRASSES: SODFARMS (AGRICULTURAL CROP USE)

For sodfarm applications, follow provisions within the **AGRICULTURAL USE REQUIREMENTS** box.

Harvesting of treated turf is prohibited until 120 hours following application.

- Do not apply more than 4 applications per year at a maximum application rate of 17.4 lb ai/A per application.
- Do not allow less than a 10 day interval between applications.

CROP	DISEASE/PEST	RATE OZ OR LBS PRODUCT	APPLICATION DIRECTIONS	RESTRICTIONS
Sod Farm	Algae	6 oz in 3 to 5 gal/1000 sq ft (16 lbs in 130 to 220 gals/A)	Begin when algae begins to appear. Repeat at 10 day intervals as long as condition persists.	Do not use on grasses grown for seed. Do not use on grasses intended for grazing, such as range or pasture grasses. Do not graze treated areas or feed clippings to livestock.
	Copper Spot Fusarium Blight <i>(F. roseum)</i> Red Thread Slime Molds	4 to 8 oz in 3 to 5 gal/1000 sq ft (11 to 22 lbs in 130 to 220 gal/A). When conditions are unusually favorable for disease, use 6 to 8 oz/1000 sq ft (16 to 22 lbs/A) and reduce intervals to 3 to 5 days.	Begin when disease appears. Repeat at 10 day intervals as long as condition persists.	
	Gray Leaf Spot <i>(Pyricularia grisea)</i>	8 oz in 3 to 5 gal/1000 sq ft (22 lbs in 130 to 220 gal/A)	Begin at first sign of disease; apply at 10 day intervals or more often during favorable disease conditions.	
	Dollar Spot <i>(Sclerotinia)</i>	6 to 8 oz in 3 to 5 gal/1000 sq ft (16 to 22 lbs in 130 to 220 gal/A) When conditions are unusually favorable for disease, use 6 to 8 oz/1000 sq ft (16 to 22 lbs/A) and reduce intervals to 3 to 5 days.	Begin when grass greens up in spring/10 to 14 days.	

(continued)

**GRASSES: SODFARMS (AGRICULTURAL CROP USE) (continued)**

CROP	DISEASE/PEST	RATE OZ OR LBS PRODUCT	APPLICATION DIRECTIONS	RESTRICTIONS
Sod Farm	Pink (Fusarium) Snow Mold	6 to 8 oz in 3 to 5 gal/1000 sq ft (16 to 22 lbs in 130 to 220 gal/A)  When conditions are unusually favorable for disease, use 6 to 8 oz/1000 sq ft (16 to 22 lbs/A) and reduce intervals to 3 to 5 days.	Apply at 2 to 6 week intervals during winter.	Do not use on grasses grown for seed.  Do not use on grasses intended for grazing, such as range or pasture grasses.  Do not graze treated areas or feed clippings to livestock.
	Leaf Spot ( <i>Helminthosporium</i> spp.) Rhizoctonia solani Brown Patch	4 oz in 3 to 5 gals/1000 sq ft (11 lbs in 130 to 220 gals/A)	Begin when disease appears. Repeat at 10-day intervals as long as condition persists.	
	Pythium Blight	8 oz in 3 to 5 gal/1000 sq ft (22 lbs in 130 to 220 gal/A)	Repeat at 5 day intervals, or more frequently if conditions are favorable for disease development.	
	Leaf Rust Stem Rust Stripe Rust	4 oz in 3 to 5 gals/1000 sq ft (11 lbs in 130 to 220 gals/A)	Begin when disease threatens. Repeat at 10 day intervals as long as disease persists.	

## SEED TREATMENTS

For commercial treatment only. Do not treat seed at or immediately before planting. Users making commercial seed applications must follow provisions within the **NON-AGRICULTURAL USE REQUIREMENTS** Box. Users conducting seed treatments on agricultural establishments must follow provisions within the **AGRICULTURAL USE REQUIREMENTS** Box.

Seed to be treated must be clean and well-cured prior to treatment.

MANZATE PRO-STICK must be applied to dry seed with conventional slurry or mist seed treating equipment, or as a plant-box application. For best results, seed must be covered uniformly with fungicide.

### Seed Bag Label Requirements

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements.

- This seed has been treated with MANZATE PRO-STICK, a fungicide containing Mancozeb.
- Do not use treated seed for feed, food, or oil purposes.

The US Environmental Protection Agency requires the following statements on containers containing seed treated with Mancozeb:

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets, or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting (such as in row ends).
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol product if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

*(continued)*



## SEED TREATMENTS *(continued)*

### REQUIRED DYE STATEMENT

Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155 (c).

When opening this bag or loading/pouring the treated seed, wear long-sleeved shirt, long pants, shoes, socks, chemical-resistant gloves, and a minimum of a NIOSH-approved particulate filtering facepiece respirator N, R, or P filter, NIOSH-approved prefix TC-84A.

After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the soil/media subsurface. Excess treated seed may be used for ethanol production only if (1) byproducts are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol byproducts that are used in agronomic practice.

CROP	DISEASES	SEED TREATMENT RATE-APPLY AS A SLURRY	
		OZ/BU	OZ/100 LBS
Barley	Bunt, Covered Smut, Damping-Off, Fake Loose Smut, Seed Decay, Seedling Blights	1.3 to 2.0	2.7 to 4.2
Corn	Damping-Off, Seed Rot, Seedling Blights	1.5 to 3.0	2.7 to 5.4
Cotton Acid Delinted	Damping-Off, Seedling Blights	--	3.0
Cotton Reginned	Damping-Off, Seedling Blights	--	6.0
Flax	Seed Decay, Seedling Blights, Damping-Off	2.0 to 4.0	3.6 to 7.1
Oat	Damping-Off, Seedling Blights, Seed Decay, Smuts	1.3 to 2.0	4.0 to 6.3
Peanut (Shelled)	Damping-Off, Seed Rots, Seedling Blights	2.0 to 4.0	8.0 to 16.0

*(continued)*

## SEED TREATMENTS *(continued)*

CROP	DISEASES	SEED TREATMENT RATE-APPLY AS A SLURRY	
		OZ/BU	OZ/100 LBS
Rice	Achyla, Other Soil and Seedborne Fungi Causing Seed Rot and Reduced Seedling Vigor	--	2.0 to 4.0 Apply before, during or after soaking in water
Rye	Bunt, Covered Smut, Damping-Off, Seed Decay, Seedling Blights	1.3 to 2.0	2.3 to 3.6
Safflower	<i>Puccinia carthami</i> (Which Causes Foot-and-Rot Disease and Foliage Rust Disease)	--	2.0
Sorghum	Covered Kernel Smut, Damping-Off, Seedling Blights, Seed Rots	1.5 to 2.5	2.7 to 4.5
Tomato	Damping-Off, Seedling Blights, Seed Rots	--	8.0
Wheat (including Triticale)	Bunt, Covered Smut, Damping-Off, Seed Decay, Seedling Blights	1.3 to 2.0	2.2 to 3.3

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** **Important**-Never allow MANZATE PRO-STICK to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of MANZATE PRO-STICK as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed in a secure dry area.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**ATTENTION:** This product contains mancozeb and ETU, chemicals known to the State of California to cause cancer in laboratory animals. ETU is also known to the State of California to cause birth defects or other reproductive harm in laboratory animals.

**IMPORTANT INFORMATION READ BEFORE USING PRODUCT**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

UPL, the UPL logo, MANZATE and PRO-STICK are trademarks of a UPL Corporation Limited Group Company.

©2023 UPL Corporation Limited Group Company

2180182-B (0823)

ESL010722-11032-083123

## NOTES

## NOTES

